

## To the fauna of the free-living nematodes (Nematoda) of high-mountain water bodies of the Western Himalayas and the Pamir

## К фауне свободноживущих нематод (Nematoda) высокогорных водоёмов Западных Гималаев и Памира

S.YA. TSALOLIKHIN

С.Я. ЦАЛОЛИХИН

*S.Ya. Tsalolikhin, Zoological Institute, Russian Academy of Sciences, 1 Universitetskaya Emb., St Petersburg 199134, Russia, E-mail: nematode@zin.ru*

Seventeen species of the nematodes, typical for the Palaearctic fauna, from the orders Tobrilida, Dorylaimida, Mononchida, Monhysterida, Chromadorida and Plectida were recorded for the first time in water bodies of the Western Himalayas and the Pamir.

В водоёмах Западных Гималаев и Памира было обнаружено 17 характерных для палеарктической фауны видов нематод из отрядов Tobrilida, Dorylaimida, Mononchida, Monhysterida, Chromadorida and Plectida.

**Key words:** free-living nematodes, Himalayas, Pamir, biogeography, new records

**Ключевые слова:** свободноживущие нематоды, Гималаи, Памир, биogeография, новые находки

### INTRODUCTION

The free-living freshwater nematofauna of mountain lakes situated over 3.500 m have been very poorly studied (Tsalolikhin, 1998). Actually, only three articles were dedicated to the freshwater nematodes of highlands (Tsalolikhin, 1998; Andrásy, 1978; Steiner, 1920). The situation is somewhat better with the alpine nematode fauna of mosses and lichens (Gadea, 1961, 1965; Micoletzky, 1929; Zullini, 1973, etc.). In high mountain lakes of the Eastern Himalayas (Nepal) 26 species of the free-living nematodes have been found, including a new species (Andrásy, 1978; Tsalolikhin, 1983). A species composition of the nematodes of the Pamir mountain range is closely related to the same of the Himalayas. The Western Himalayan Biogeographic Expedition of the St Petersburg Association of Scientists and Scholars under guidance of Dr. L.Ja. Borkin (20 April – 14 May 2013) collected additional information on alpine

water bodies of the Himalayas, and the International Pamir Biological Expedition granted by the National Geographic Society (July – August 2013) added new data on the nematode fauna of the Pamir.

### MATERIAL AND METHODS

Benthos samples were taken from six water bodies in the Western Himalayas and from seven ones in the Central Pamir. A distribution of the nematodes in water bodies is presented in Table 1.

### List of water bodies where nematodes were found

The Western Himalayas (Jammu and Kashmir State, India):

- 1) Mansar Lake: 665 m; 75.151°E, 32.696°N (22.IV.2013);
- 2) Wullar Lake: 1506 m; 74.56° E, 34.378°N (1.V.2013);

- 3) Tsomo Riri Lake: 4469 m; 78.15°E, 33.768°N (10.V.2013);  
 4) Nubra River: 3152 m; 77.54°E, 34.777°N (7.V.2013);  
 5) Spring: 3148 m; 77.555°E, 34.758°N (7.V.2013);  
 6) Termal spring: 4381 m; 78.315°E, 33.223°N (10.V.2013).  
The Pamir (Gorno-Badakhshan Autonomous Region, Tajikistan):  
 7) Rangkul Lake: 3752 m; 74.264°E, 38.471°N (30.VII.2013);  
 8) Karakul Lake: 3883 m; 73.554°E, 39.017°N (2.VIII.2013);  
 9) Khargush Lake: 4213 m; 73.068°E, 37.431°N (21.VII.2013);  
 10) Ail-Utek spring: 3659 m; 73.04°E, 38.558°N (31.VII.2013);  
 11) Issyk-bulok termal spring: 3720 m; 72.977°E, 37.771°N (24.VII.2013);  
 12) Kokuybel puddle: 3595 m; 73.067°E, 38.506°N (1.VIII.2013);  
 13) Karasu puddle: 3749 m; 73.973°E, 38.176°N (27.VII.2013).

**Table 1.** Distribution of freshwater nematodes in the Western Himalayas and the Pamir.

Species	Water bodies (listed in "Material and methods" section)												
	West Himalayas						Central Pamir						
	1	2	3	4	5	6	7	8	9	10	11	12	13
<b>Tobrilida</b>													
<i>Tobrilus gracilis</i> (Bast., 1865)			+										+
<i>T. helveticus</i> (Hofmaenner, 1914)							+						
<i>Brevitobrilus vibratus</i> (Sukul, 1967)	+												
<i>Neotobrilus tantloji</i> (Sukul, 1971)												+	
<i>Tripyla glomerans</i> Bast., 1865		+											
<b>Dorylaimida</b>													
<i>Dorylaimus fodori</i> Andrassy, 1988							+		+			+	
<i>Crocodylaimus borchuk</i> Tsalolikhin, 2014				+		+							
<i>Laimydorus</i> sp.								+					
<i>Labronema ferox</i> Thorne, 1939							+						
<b>Mononchida</b>													
<i>Mononchus truncatus</i> Bast., 1865													+
<i>Mononchus</i> sp.		+											
<b>Monhysterida</b>													
<i>Monhystera wangi</i> Wu et Hoeppl, 1929	+		+		+								
<i>Eumonhystera similis</i> (Buetschli, 1873)				+									
<b>Chromadorida</b>													
<i>Achromadora terricola</i> (de Man, 1880)													+
<i>Prodesmodora</i> sp.					+								
<b>Plectida</b>													
<i>Plectus palustris</i> de Man, 1880								+					+
<i>Plectus rhizophilus</i> de Man, 1880				+									

The following abbreviations, except for most commonly accepted ones, are used in the descriptions of the species and in tables: cl-I – a distance between the cloaca and the first supplement; Q – the length of the gonad; Sp – a spicule length, SR – the length of the supplement row; Sp/cl-I – a ratio of a spicule length to a distance between the cloaca and the first supplement; Sp/SR – a ratio of a spicule length to the length of the supplement row; SR/L – a ratio of the length of the supplement row to a body length; Suppl/SR – a ratio of a distance between the supplements to the length of the supplement row and V-an/cd – a ratio of a distance between the vulva and the anus to a tail length. Roman numerals in the last five columns of Table 4 denote a ratio of a distance between the supplements to the length of supplement row, in percentage terms. All measurements were made in micrometers.

## RESULTS

Brief descriptions of some notable species are given below.

### *Tobrilus gracilis* (Bastian, 1865) (Figs 1–6)

*Description. Females* (n = 8): L = 1519–1891 (1710±47) µm, a = 23.4–30.5 (27±1.2), b = 3.9–5.4 (4.7±0.15), c = 7.4–9.6 (8.6±0.3), c' = 5–7, V = 41–51 (45±1)%. Head 28–30 (29±0.5) µm wide, cephalic setae 10–16 (13±1) µm long, buccal cavity 9–10 × 11–13 µm, total depth of stoma 21–28 (25±1) µm. Oesophagus 320–440 (364±15) µm long, NR = 30%. Female reproductive system: Q<sub>1</sub> = 170–245 µm, Q<sub>2</sub> = 210–265 µm; V-an/cd = 3.4–4.4 (3.7±0.1). Rectum 35–49 (45±2) µm long, tail 170–245 (200±8) µm long, subterminal seta absent.

*Males* (n = 6): L = 1430–1883 (1610±77) µm, a = 28.3–38.4 (32.9±2), b = 4.3–5.8 (5.1±0.3), c = 11.6–17.4 (13.5±1), c' = 3–4. Head 22–26 (24±0.5) µm wide, cephalic setae 11–12 µm long, buccal cavity 7–9 × 10–11 µm, total depth of stoma 19–21 µm.

Oesophagus 265–363 (320±21) µm long, NR = 33%. Tail 108–130 (120±4) µm long, subterminal seta absent.

*Notes.* The Himalayan *T. gracilis* is somewhat smaller compared with “typical” specimens of the species, their parametric data almost completely coincide (Tables 2, 3), particularly for the supplementary apparatus (Tsalolikhin, 1998).

### *Tobrilus helveticus* (Hofmaenner, 1914)

*Description. Females* (n = 3): 1690–1825 (1777) µm, a = 26–30.3 (28.2), b = 4.9–5.7 (5.3), c = 7.7–9.2 (8.3), c' = 5.4–6.3 (6), V = 46–49 (48)%. Head 20–25 (23) µm wide, cephalic setae 5 µm long, buccal cavity 6–9 (7) × 9 µm, total depth of stoma 19–23 (21) µm. Oesophagus 294–370 (335) µm, NR = 32–35 (33)%. Female reproductive system underdeveloped; V-an/cd = 3–4. Rectum 29–36 (33) µm long, tail 196–220 (212) µm long, subterminal seta absent.

*Notes.* The specimens from the Pamir mainly differ from “typical” *T. helveticus* in a slightly shorter tail. Several specimens of *T. helveticus* with short tails were found in Po River, Italy (Zullini, 1974) and in Volga River (Gagarin, 1993).

### *Brevitobrilus vibratus* (Sukul, 1967) (Figs 7–12)

*Description. Females* (n = 2): 990–1137 µm, a = 20.7–29.1, b = 5.1–5.4, c = 6.7–7.1, c' = 7–8, V = 43–45%. Head 17–23 µm wide, cephalic setae 4 µm long, buccal cavity 8–9 × 11–13 µm, total depth of stoma 22–28 µm, distance between tops of onchia 6 µm. Oesophagus 196–212 µm, NR = 39–49%. Females gonads two, ovaries reflexed: Q<sub>1</sub> = 98–136 µm, Q<sub>2</sub> = 98–155 µm, eggs 45 × 23–30 µm; V-an/cd = 3. Rectum 26–30 µm long, tail 147–159 µm long, subterminal setae absent.

*Male:* L = 1194 µm, a = 44, b = 5.8, c = 11.1, c' = 4, suppl. 6. Head 21 µm wide, cephalic setae 5 µm long. Oesophagus 204 µm long, tail 108 µm long, subterminal setae absent.

*Notes.* The rare species, originally described from India (Sucul, 1967), is morphologically very similar to *B. stefanskii* (Micoletzky 1925), and it was the reason for a synonymisation of these two names (Tsalolikhin, 2001). The differences between these two species consist in sizes of the spicules: 26  $\mu\text{m}$  vs 30–40  $\mu\text{m}$  and in the index sp/cl-I: 124 vs 113, respectively.

Both species belong to the *stefanskii*-group (group A) of the genus *Brevitobrilus* Tsalolikhin, 2001. Apart from India, *B. vibratus* was also found in Mongolia (Tsalolikhin, 1985). Comparative measurements of the supplementary apparatus of *B. vibratus* and *B. stefanskii* are presented in Table 4. Data for *B. stefanskii* are given by Micoletzky (1925) and Tsalolikhin (2001).

**Table 2.** Comparative measurements of *Tobrilus gracilis*.

Characters	Data for species by Tsalolikhin, 2009	Specimens from Tso Moriri Lake
Females		(n = 8)
L	1670–3300 (2185)	1519–1891 (1710 $\pm$ 47)
<i>a</i>	19.6–40 (25.5)	23.4–30.5 (27 $\pm$ 1.2)
<i>b</i>	4.4–6.7 (5.4)	3.9–5.4 (4.7 $\pm$ 0.15)
<i>c</i>	7–15.4 (9.9)	7.4–9.6 (8.6 $\pm$ 0.3)
V%	40–54 (45)	41–51 (45)
<i>c'</i>	4–5 (4.7)	5–7.5 (6.5)
NR%	27–34 (31)	30
Oesophagus	339–530 (425)	320–440 (364 $\pm$ 15)
Tail	190–283 (218)	170–245 (200 $\pm$ 8)
V-an/cd	(4.4)	3.4–4.4 (3.7 $\pm$ 0.1)
Q <sub>1</sub>	(280)	170–245 (220)
Q <sub>2</sub>	(280)	210–280 (227)
Rectum	(53)	35–49 (45 $\pm$ 2)
Head	31–40 (35)	28–30 (30 $\pm$ 1)
Cephalic setae	10–13 (12)	10–16 (13 $\pm$ 1)
Cephalic setae/head,%	30–40 (37)	45
Buccal cavity (width)	11–13 (12)	9–10
Stoma (depth)	23–28 (25)	21–28 (25 $\pm$ 1)
Males		(n = 6)
L	1460–2600 (1954)	1430–1883 (1610 $\pm$ 77)
<i>a</i>	20–45 (31)	28.3–38.4 (32.9 $\pm$ 2)
<i>b</i>	4.3–6.6 (5.4)	4.3–5.8 (5.1 $\pm$ 0.3)
<i>c</i>	10.4–17 (14)	11.6–17.4 (13.5 $\pm$ 1)
NR%	31–34 (33)	33
Oesophagus	330–413 (370)	265–363 (320 $\pm$ 21)
Trophico-genital part	1133–1716 (1438)	1030–1412 (1170 $\pm$ 68)
Tail	120–160 (146)	108–130 (120 $\pm$ 4)

**Table 3.** Measurements of supplementary apparatus of *Tobriulus gracilis*.

Suppl/SR	Data for species by Tsalolikhin, 2009	Specimens from Tso Moriri Lake
Cl-I	15–30 (20)	17–22 (20)
I-II	12–17 (14)	15–17 (16)
II-III	8–12 (10)	9–12 (11)
III-IV	6–12 (9)	7–10 (9)
IV-V	4–10 (8)	9–11 (9)
V-VI	6–10 (8)	9–12 (11)
VI-VII	10–16 (13)	9–12 (10)
VII-VIII	15–20 (17)	11–21 (14)
Sp/cl-I, %	70–80 (74)	66–83 (79)
SR/L, %	12–20 (17)	11–17 (14)
Sp/SR, %	13–19 (15)	9–16 (15)
Sp, $\mu\text{m}$	36–40 (38)	28–37 (33)
SR, $\mu\text{m}$	256–397 (328)	207–245 (221)

***Neotobriulus tantloji*** (Sukul, 1971)  
(Figs 13–15)

*Description.* *Juvenile:* 920  $\mu\text{m}$ ,  $a = 23$ ,  $b = 4.7$ ,  $c = 10.6$ . Head 16  $\mu\text{m}$  wide, cephalic setae 4  $\mu\text{m}$  long, buccal cavity  $4 \times 4 \mu\text{m}$ , total depth of stoma 18  $\mu\text{m}$ , distance between tops of onchia 6  $\mu\text{m}$ . NR = 40%.

*Notes.* This very rare species was found only once in a thermal spring in India. One female and three juveniles were described by Sukul, (1971). The author listed the following characters of larvae: L = 760 – 940  $\mu\text{m}$ ,  $a = 22.8 - 26.4$ ,  $b = 4.2 - 4.4$ ,  $c = 6.8 - 7.7$ . The measurements of the juvenile from the Pamir corresponds these date, but Pamirian specimen has the shorter tail. It is

noteworthy that this specimen was found in the thermal spring as well as the type specimens of *N. tantloji*.

***Tripyla glomerans*** Bastian, 1865

*Notes.* The species from Wuler Lake in the Western Himalayas is presented only by juveniles. The range of the species is extremely wide but does not extend beyond the Eastern Hemisphere; it can be reliably regarded as the Palaearctic species (Tsalolikhin, 1983). It has been previously recorded for the Himalayas (Andrássy, 1978).

***Dorylaimus fodori*** Andrásy, 1988

*Notes.* The species was described from paddy field in Bangalor State, India (Andrássy, 1988). The finding in the Pamir extended the range of the species and indicated a wide distribution of it in Asia. The main morphometric characters of the species are presented in Table 5.

***Crocodyrlaimus borchuk***  
Tsalolikhin, 2014


*Notes.* The species was described from the Nubra River, the Western Himalayas (Tsalolikhin, 2014b). The main morphometric characters of the species are presented in Table 6.

***Laimydorus*** sp.

*Description.* *Female:* L = 1886  $\mu\text{m}$ ,  $a = 33.1$ ,  $b = 4.7$ ,  $c = 11.6$ ,  $c' = 5.6$ ,  $V = 45\%$ . Head 17  $\mu\text{m}$  wide; spear 23  $\mu\text{m}$  long. Oesophagus 402  $\mu\text{m}$  long; V-an/cd = 5.4,  $Q_1 = 282 \mu\text{m}$ ,  $Q_2 = 302 \mu\text{m}$ . Rectum 49  $\mu\text{m}$  long, prerectum 74  $\mu\text{m}$  long. Tail 163  $\mu\text{m}$  long.


**Table 4.** Measurements of the supplementary apparatus of *Brevitobriulus vibratus* and *B. stefanskii*.

Species	Sp, $\mu\text{m}$	SR, $\mu\text{m}$	SR/L	Sp/SR	Sp/cl-I	cl-I	I-II	II-III	III-IV	IV-V	V-VI
<i>B. vibratus</i>	26	180	15	14	63	12	23	15	17	15	17
<i>B. stefanskii</i>	35	220	17	16	87	18	20	15	15	15	16



**Figs 1–6.** *Tobrilus gracilis*. 1, entire body of female; 2, entire body of male; 3, head, dorsal view; 4, head, lateral view; 5, tail of female; 6, posterior end of male body.





**Figs 7–15.** Nematodes of genera *Brevitobrilus* and *Neotobrilus*. 7–12, *Brevitobrilus vibratus*. 13–15, *Neotobrilus tantloyi*. 7, entire body of female; 8, head, dorsal view; 9, 13, head, lateral view; 10, vulvar section; 11, 15, tail of female; 12, posterior end of male body; 14, oesophagus.

**Table 5.** Measurements of *Dorylaimus fodori*.

Characters	Males (n=6)	Females (n=5)
L	3561–4310 (3842±126)	4130–4940 (4540±167)
Oesophagus	796–922 (858±23)	825–970 (897±31)
Tail	35–47 (42±2)	214–291 (260±19)
Prerectum+rectum	388–485 (455±19)	337–355 (345±9)
Spicula	74–97 (82±4)	–
Supplements	44	–
Head	18–19	18–19
Spear	43 × 5	45 × 5
<i>a</i>	34.5–53.2 (42.7±1.4)	33.2–50.9 (45.9±4.3)
<i>b</i>	4.2–4.9 (4.5±0.1)	4.7–5.4 (5.1±0.1)
<i>c</i>	75.8–110 (92.1±7.2)	9.3–16.9 (15.1±2)
<i>c'</i>	0.7–1 (0.9±0.05)	5.4–7.1 (5.9±0.4)
V%	–	41–46 (43±1)
V-an/cd	–	8.3–9.4 (8.8±0.3)

**Table 6.** Measurements of *Crocodyrlaimus borchuk*.

Characters	Males (n=4)	Females (n=4)
L	1560–1720 (1654)	1582–1883 (1776)
Oesophagus	294–345 (314)	279–326 (306)
Tail	61–82 (72)	120–136 (129)
Prerectum	115–120 (118)	80–90 (87)
Rectum	30–40 (35)	32–36 (34)
Spicula	35–38 (37)	–
Supplements	18	–
SR	65–68 (67)	–
Head	10–11	10–11
Spear	14–16 (15)	14–16 (15)
Vagina	–	24–27 (26)
Q <sub>1</sub>	–	250–340
Q <sub>2</sub>	–	250–340
Egg	–	65–80 × 26–27
<i>a</i>	42–48.4 (45.5)	38.6–45.1 (43.3)
<i>b</i>	5.0–5.6 (5.3)	5.4–6.2 (5.8)
<i>c</i>	61–82 (72)	13.3–15.7 (14.0)
V%	–	43–48 (45)
<i>c'</i>	1	6.3–7.8 (7.2)
V-an/cd	–	6–8
NR%	32–37 (35)	36



***Labronema ferox*** Thorne, 1939

*Description. Females* (n = 2): L = 2207–2782  $\mu\text{m}$ ,  $a = 22.7\text{--}25.3$ ,  $b = 3.6\text{--}3.9$ ,  $c = 63\text{--}80$ ,  $c' = 0.8$ ,  $V = 48\text{--}50\%$ . Head 24–25  $\mu\text{m}$  wide; spear 35–36  $\times$  5  $\mu\text{m}$ , aperture of spear 13  $\mu\text{m}$  (1/3 length of spear); guiding ring doubl. Oesophagus 620–704  $\mu\text{m}$ ; NR = 45%. Female reproductive system underdeveloped, depth of vagina 50  $\mu\text{m}$ , V-an/cd = 31–40. Rectum 65–68  $\mu\text{m}$  long, prerectum 100–110  $\mu\text{m}$  long. Tail 35  $\mu\text{m}$  long.

*Notes.* The species was described from the U.S.A. (Thorne, 1939) but was also recorded for the Himalayas (Zullini, 1973). In addition, *L. loeffleri* Andr ssy, 1978, a species close related with *L. ferox*, was found in the Himalayas in the Western Lobuche Lake at an altitude of 5180 m.

***Monhystera wangi*** Wu et Hoepfly, 1929

*Description. Females* (n = 8): 742–930 (856 $\pm$ 22)  $\mu\text{m}$ ,  $a = 28.2\text{--}32.3$ ,  $b = 5.4\text{--}7.4$  (6.5 $\pm$ 0.2),  $c = 4.9\text{--}5.7$  (5.2 $\pm$ 0.1),  $c' = 7\text{--}11$  (9 $\pm$ 0.5),  $V = 55\text{--}62$  (59 $\pm$ 1)%.

*Male:* L = 781  $\mu\text{m}$ ,  $a = 33.9$ ,  $b = 6.2$ ,  $c = 5.8$ ,  $c' = 7$ , spic. 30  $\mu\text{m}$ , gub. 6  $\mu\text{m}$ . Head 10–12  $\mu\text{m}$  wide, cephalic setae very delicate, 3–4  $\mu\text{m}$  long, stoma almost funnel-shaped. Center of amphid situated at 7–8  $\mu\text{m}$  from the anterior body end, diameter of amphid 4  $\mu\text{m}$ . Oesophagus cylindrical but widening towards the base (especially in male), 114–146 (132 $\pm$ 3)  $\mu\text{m}$  long; oesophago-intestinal glands large. NR = 50–54%. Female reproductive system monodelphic, 320–350  $\mu\text{m}$  long, egg 45–50  $\times$  10–21  $\mu\text{m}$ , depth of vagina 23  $\mu\text{m}$ ; V-an/cd = 1–1.2. Rectum 16–23 (19 $\pm$ 1)  $\mu\text{m}$  long. Female tail 140–182 (164 $\pm$ 5)  $\mu\text{m}$  long, male tail 135  $\mu\text{m}$  long. Paracloacal cuticle plicated.

*Notes.* The species was described from fresh water bodies in South China (Wu & Hoepfly, 1929) and recorded for Mongolia (Tsalolikhin, 1985) and the Pamir (Tsalolikhin, 1998).

***Achromadora terricola*** (de Man, 1880)

*Description. Female:* L = 926  $\mu\text{m}$ ,  $a = 25.7$ ,  $b = 6.3$ ,  $c = 8$ ,  $c' = 5.3$ ,  $V = 48\%$ . Head

18  $\mu\text{m}$  wide, cephalic setae 5  $\mu\text{m}$  long, center of amphid situated at 14  $\mu\text{m}$  from the anterior body end, diameter of amphid 5  $\mu\text{m}$ . Oesophagus 147  $\mu\text{m}$  long, bulbus 28  $\times$  24  $\mu\text{m}$ . V-an/cd = 3.1. Rectum 48  $\mu\text{m}$  long. Tail 117  $\mu\text{m}$  long.

*Notes.* This widely distributed species was previously twice recorded for the Himalayas (Andr ssy, 1978; Zullini, 1973).

***Prodesmodora*** sp.

*Description. Female:* L = 857  $\mu\text{m}$ ,  $a = 26.8$ ,  $b = 7.2$ ,  $c = 5.7$ ,  $c' = 8.8$ ,  $V = 42\%$ . Head 19  $\mu\text{m}$  wide, cephalic setae very thin, about 3  $\mu\text{m}$  long, center of amphid situated at 18  $\mu\text{m}$  from the anterior body end, diameter of amphid about 4  $\mu\text{m}$ . Oesophagus 119  $\mu\text{m}$  long, bulbus 23  $\times$  19  $\mu\text{m}$ . NR = 57%. Female reproduction system didelphic:  $Q_1 = 146$   $\mu\text{m}$  (egg 57  $\times$  23  $\mu\text{m}$ ),  $Q_2 = 126$   $\mu\text{m}$  (egg 58  $\times$  23  $\mu\text{m}$ ); V-an/cd = 2.3. Rectum 31  $\mu\text{m}$  long, tail 150  $\mu\text{m}$  long, anal body diameter 17  $\mu\text{m}$ .

*Notes.* In this female the length of the rectum is nearly twice as long as the anal diameter. The long rectum within the genus is only in *P. terricola* Altherr, 1952 and *P. minuta* Schneider, 1937, however, in the first species the rectum length is 3–4 times as an anal body diameter, whereas in the last species a body length is much shorter than in the Himalayan specimen. In *P. minuta*, the rectum is either shorter or less than 1.5 times as an anal body diameter. This character allows me to consider the described specimen as a new species; however, the single specimen is not sufficient for naming and describing a new species. The genus *Prodesmodora* Micoletzky, 1923 is widely distributed around the world. *P. circulata* (Micoletzky, 1913) is most frequently found in fresh (and brackish) water bodies but reliability of identification of this species is questionable in some cases (Tsalolikhin, 2014a).

Other species included in Table 1, were collected as single specimens and usually as larvae. Species such as *Tripyla glomerans* Bastian, 1865, *Mononchus truncatus* Bas-

tian, 1865, and *Eumonhystera similis* (Bütschli, 1873) have been previously found in the Himalayas. They are universally distributed or belong to the Palearctic, Ethiopian and Sino-Indian Regions (Tsalolikhin, 1983).

## ACKNOWLEDGEMENTS

I am very grateful to Leo Borkin (St Petersburg) and Spartak Litvinchuk (St Petersburg) for the valuable material used in the paper.

## REFERENCES

- Andrássy I.** 1978. Fresh-water Nematodes from the Himalayas (Nepal). *Opuscula Zoologica Budapest*, **15**(1–2): 3–21.
- Andrássy I.** 1988. The superfamily Dorylaimoidea – a review. Family Dorylaimidae. *Opuscula Zoologica Budapest*, **23**: 3–63.
- Gadea E.** 1961. Nota sobre algunos nematodos muscícolas del Nepal. *Miscel-lània Zoològica*, **1**(4): 3–7.
- Gadea E.** 1965. Nematodos muscícolas de los Andes del Perú. *Miscel-lània Zoològica*, **2**(1): 3–12.
- Gagarin W.G.** 1993. *Svobodnozhivushchie nematody presnykh vod Rossii i sopredel'nykh stran (otryady Monhysterida, Araeolaimida, Chromadorida, Enoplida, Mononchida)* [Free-living nematodes in freshwater-bodies of Russia and adjacent countries (orders Monhysterida, Araeolaimida, Chromadorida, Enoplida, Mononchida)]. St Petersburg: Gidrometeoizdat. 350 p. (In Russian).
- Micoletzky H.** 1929. Zoologische Ergebnisse der Deutsch-Russischen Alai-Pamir-Expedition. *Zoologischer Anzeiger*, **84**: 244–252.
- Steiner G.** 1920. Freilebende Süßwassernematoden aus peruanischen Hochgebirgsseen. *Revue suisse de Zoologie*, **28**(2): 11–44.
- Sukul N.C.** 1967. A new species of *Tobrilus* with notes on its biology. *Indian Journal of Helminthology*, **19**: 113–117.
- Sukul N.C.** 1971. New free-living nematodes from the thermal spring of Tantloya, Bihar. *Indian Journal of Helminthology*, **23**: 135–145.
- Thorne G.** 1939. A monograph of the Nematodes of the superfamily Dorylaimoidea. *Capita Zoologica*, **8**: 3–258.
- Tsalolikhin S.Ya.** 1983. *Nematody semeystv Tobrilidae i Tripylidae mirovoy fauny* [Nematodes from the families Tobrilidae and Tripylidae of the world fauna]. Leningrad: Nauka. 232 p. (In Russian).
- Tsalolikhin S.Ya.** 1985. *Nematody presnykh i solonovatykh vod Mongolii* [Nematodes of fresh and brackish waters of Mongolia]. Leningrad: Nauka. 115 p. (In Russian).
- Tsalolikhin S.Ya.** 1998. On nematode fauna of high mountain lakes in the Pamir and Himalayas. *Zoologicheskii Zhurnal*, **77**(9): 1058–1063. (In Russian).
- Tsalolikhin S.Ya.** 2001. Notes on composition of the genus *Brevitobrilus* Tsalolikhin, 1981 (Nematoda, Enoplida: Tobrilidae). *Zoosystematica Rossica*, **9**(1): 25–35.
- Tsalolikhin S.Ya.** 2009. A review of the genus *Tobrilus* (Nematoda, Enoplida: Tobrilidae): “Classical species”. *Zoologicheskii Zhurnal*, **88**(7): 783–793. (In Russian).
- Tsalolikhin S.Ya.** 2014a. Review of the *Prodesmodora* (Nematoda, Desmodorida). *Zoosystematica Rossica*, **23**(1): 3–6.
- Tsalolikhin S.Ya.** 2014b. A review of the genus *Crocodylaimus* (Nematoda: Dorylaimida). *Trudy Zoologicheskogo Instituta Rossiyskoy Akademii Nauk* [Proceedings of the Zoological Institute of the Russian Academy of Sciences], **318**: 280–287. (In Russian).
- Wu H. & Hoeppl R.** 1929. Free-living nematodes from Fookien and Chekiang. *Archiv für Tropenhygiene*, **33**: 35–43.
- Zullini A.** 1973. Su alcuni Nematodi di alta quota del Nepal. *Khumbu Himal*, **4**: 401–412.
- Zullini A.** 1974. The nematodical population of the Po River. *Bollettino di Zoologia*, **41**: 183–210.

Received September 15, 2014 / Accepted October 24, 2014

Editorial responsibility: D.A. Gapon, A.Yu. Ryss